



Glaciation and deglaciation of the Stump Cross area, Yorkshire Dales, northern England, determined by terrestrial cosmogenic nuclide (^{10}Be) dating.

Wilson, P., Lord, T. C., & Rodes, A. (2017). Glaciation and deglaciation of the Stump Cross area, Yorkshire Dales, northern England, determined by terrestrial cosmogenic nuclide (^{10}Be) dating. *Cave and Karst Science*, 44(2), 76-81.

[Link to publication record in Ulster University Research Portal](#)

Published in:
Cave and Karst Science

Publication Status:
Published (in print/issue): 06/10/2017

Document Version
Publisher's PDF, also known as Version of record

General rights
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.



Glaciation and deglaciation age of the Stump Cross area, Yorkshire Dales, northern England, determined by terrestrial cosmogenic nuclide (^{10}Be) dating

Peter WILSON ¹, Tom C LORD ² and Ángel RODÉS ³

¹ Environmental Sciences Research Institute, School of Geography and Environmental Sciences, Ulster University, Coleraine, Co. Londonderry, BT52 1SA, UK.

² Lower Winskill, Langcliffe, Settle BD24 9PZ, UK, and Department of Archaeology, University of York, King's Manor, York, YO1 7EP, UK.

³ NERC Cosmogenic Isotope Analysis Facility, Scottish Universities Environmental Research Centre, Rankin Avenue, East Kilbride, G75 0QF, UK.

Corresponding author: Peter Wilson (*E-mail*: p.wilson@ulster.ac.uk)

Abstract: Terrestrial cosmogenic nuclide (^{10}Be) surface-exposure ages are reported for three glacially-transported gritstone boulders and one glacially-scoured exposure of gritstone bedrock in the vicinity of Stump Cross Caverns, North Yorkshire. Although the ages do not form a statistically consistent cluster, three of them nevertheless indicate that the transport and deposition of boulders was by ice of the last (Late Devensian) glaciation. The ages provide evidence for glacier ice at the Wharfe–Nidd interfluvium, in contrast to previously held views that these uplands had remained above the level of the last ice sheet. The youngest of the three ages on boulders (~18.5 ka) is taken as the best estimate for deglaciation of the area. This is consistent both with surface-exposure ages from sites elsewhere around the southern margin of the Yorkshire Dales and with uranium-series dated speleothems in Stump Cross Caverns. Together these results reveal that deglaciation of the Dales was most likely well advanced by ~18–16 ka, facilitating the rejuvenation of surface and subsurface karstic processes.

Keywords: Glaciation, deglaciation, terrestrial cosmogenic nuclide dating, uranium-series dating, ice-transported boulders, speleothem.

Received: 22 March 2017; Accepted: 25 May 2017.

Refinement of the timescale regarding the last glacial/deglacial cycle for certain areas of northern England has been advanced in recent years through applications of optically stimulated luminescence (OSL) dating of loessic sediment and glacio-lacustrine/fluvial sediments (Telfer *et al.*, 2009; Bateman *et al.*, 2015), and terrestrial cosmogenic nuclide (TCN) dating of ice-transported boulders and glacially-scoured bedrock (Vincent *et al.*, 2010; Wilson *et al.*, 2013; Livingstone *et al.*, 2015). In addition, the interpretation of the geomorphological and sedimentological evidence displayed by glacial landforms and materials has provided valuable information concerning the pattern and style of glacial events (Livingstone *et al.*, 2012; Chiverrell *et al.*, 2016). Even so, questions remain over whether some parts of the region were glaciated during the Last Glacial Maximum (LGM: ~26.5–19 ka BP; Clark *et al.*, 2009) of the Late Devensian sub-stage, and for other areas there is a paucity of detailed information concerning when they emerged from beneath the ice.

In the Yorkshire Dales National Park, deglaciation of limestone terrain is constrained by ages from four sites, all of which occur close to its southern boundary and are a maximum distance of

41 km apart (Sutcliffe *et al.*, 1985; Atkinson *et al.*, 1986; Telfer *et al.*, 2009; Wilson *et al.*, 2012). The ages suggest the sites became free of ice ~18–16 ka BP (Wilson and Lord, 2014). However, this apparently consistent picture of deglaciation within a relatively narrow timeframe is undermined by the large 1σ uncertainties of 2–4 ka on two of the ages (speleothems at White Scar Cave: 17 ± 4 ka (Atkinson *et al.*, 1986), and Stump Cross: 17 ± 2 ka (Sutcliffe *et al.*, 1985)). It is therefore important that additional geochronological data are obtained in order to establish whether certain areas were or were not glaciated and, if glaciated, when deglaciation occurred, and by inference provide a maximum age for the resumption of surface and sub-surface karstic processes.

In this respect the Stump Cross area on the Wharfe–Nidd interfluvium was identified as a location about which there is limited information concerning its glacial/deglacial history and where it has even been suggested that the tills there may pre-date the Devensian stage (>115 ka; Smith 1977; Catt, 1991; Baker *et al.*, 1996). At present, dated speleothems from Stump Cross Caverns provide the only chronology for Late Devensian events in the area, with U–Th ages of 17 ± 2 ka and 15 ± 1 ka (Sutcliffe

et al., 1985) indicating the renewal of groundwater flow and calcite deposition following the removal of glacial ice cover and/or permafrost degradation after the LGM. In the vicinity of the Caverns, there are several large ice-transported boulders of coarse-grained Millstone Grit sandstone (“gritstone”). These boulders were targeted for TCN dating with the aims of establishing: (1) whether the last ice-sheet covered the area and was responsible for transporting the boulders, and (2) the timing of deglaciation and associated deposition of the boulders.

Cosmogenic nuclides are formed *in situ* when a rock surface is bombarded by high-energy neutrons and subatomic particles known as cosmic-rays. This interaction alters elements contained in the minerals of the rock and new nuclei are created (cosmogenic nuclides). The most commonly used nuclides, beryllium-10 (^{10}Be), aluminium-26 (^{26}Al) and chlorine-36 (^{36}Cl) accumulate in rock over time, depending upon the half-life of the nuclide, the rate of rock surface erosion, the composition of the rock and the intensity of the bombardment. Measurement of the concentration of a nuclide and knowing its rate of production and half-life, enables the determination of how long a rock surface has been exposed to cosmic radiation (Gosse and Phillips, 2001; Phillips, 2001; Cockburn and Summerfield, 2004). Implicit in the meaningful interpretation of surface exposure ages relating to glaciation/deglaciation is that the rock surface must have been exposed continuously and without disturbance since the loss of ice cover. Furthermore, cosmogenic nuclides that accumulated during a previous period of exposure (i.e. inherited nuclides) should not be present.

Stump Cross

The Stump Cross area (360m OD; National Grid Reference SE 089 635; Figs 1 and 2) is on the broad interfluvium between the valleys of Wharfedale, to the west, and Niddersdale, to the east, and is ~30–35km inside (i.e. north of) the LGM ice limit (Clark *et al.*, 2004). The bedrock consists of Dinantian limestones overlain to the north by rocks of the Namurian Millstone Grit Group (specifically the Grassington Grit Formation) (Dunham and Stubblefield, 1945).




Figure 1: The location of Stump Cross in relation to the maximum extent of the British–Irish Ice Sheet during the Last Glacial Maximum (after Scourse and Furze, 2001; Clark *et al.*, 2004; Sejrup *et al.*, 2005; Bradwell *et al.*, 2008; McCarroll *et al.*, 2010; Ó Cofaigh *et al.*, 2012).




Figure 2: Locations of Stump Cross Caverns, Nursery Knott, and the boulders (red dots) sampled for TCN dating. The geological boundary is based upon Dunham and Stubblefield (1945).

Sample code	Grid reference	Latitude ($^{\circ}\text{N}$)	Longitude ($^{\circ}\text{W}$)	Altitude (m OD)	Thickness (cm)	Density (g cm^{-3})	Topographic shielding
NK-01	SE 08229 63970	54.07142	1.87424	365	1.5	2.32	1.0000
NK-02A	SE 08224 63974	54.07145	1.87430	363	3.0	2.35	0.9663
NK-03	SE 07884 63783	54.06975	1.87950	360	1.5	2.38	0.9980
NK-04	SE 08255 63761	54.06954	1.87384	370	1.5	2.41	0.9993

Table 1: Details of samples for TCN dating from Nursery Knott.

Stump Cross Caverns are one of the most intensively dated cave systems in the UK, with 55 age determinations, ranging from ~225 ka to ~8 ka, pertaining to phases of flowstone development in, and faunal occupation of, the Caverns (Sutcliffe *et al.*, 1985; Baker *et al.*, 1996; Gilmour *et al.*, 2007; Caseldine *et al.*, 2008; O'Connor and Lord, 2013). In spite of this wealth of sub-surface information there is little detail concerning the geochronology of surface processes, materials and events. Both Raistrick (1931) and Tillotson (1934) inferred the former presence of glacier ice in the Stump Cross area, but their chronology was hampered by the absence of an established dating framework. In contrast, Smith (1977) thought that the local glacial drift was of Wolstonian (~300–130 ka) or earlier age, implying that the area had not been inundated by Devensian (~115–13 ka) ice, including at the LGM. O'Connor and Lord (2013) suggested that the area was not covered by warm-based ice during Quaternary cold (glacial) stages because Stump Cross Caverns lack the laminated clays that characterise glacial events in the Victoria Cave sedimentary record, 25 km to the west (Lundberg *et al.*, 2010). However, the presence of ice-transported gritstone boulders on the limestone north and west of Stump Cross indicates the former presence of south-going glacier ice, but hitherto the age of this glacial event has not been established conclusively. Even though the area is regarded as within the limit of the last ice sheet at the LGM (Clark *et al.*, 2004) it has not been proven that the high ground in this area was ice covered at that time.

Methods

Samples for TCN (^{10}Be) dating were collected from three ice-transported boulders and one exposure of coarse-grained Millstone Grit sandstone in the vicinity of Nursery Knott, ~1km northwest of Stump Cross Caverns (Fig.2, Table 1). Sample NK-01 was from the top of a 2m-high boulder (Figs 3A and B) that appears to have been pushed or dragged southwards by about two metres, presumably by glacier ice, from the underlying bedrock that now forms a plinth immediately north the boulder. The plinth stands <0.5m above the surrounding vegetated ground and the boulder rests partly on its southern edge; sample NK-02A was obtained from the centre of the plinth (Figs 3C and D). The other two samples (NK-03 and -04) were from the uppermost surfaces of 1m-high boulders that had been transported minimum distances of ~200m from the gritstone outcrop to their current positions on the limestone (Figs 3E and F). Samples were collected from these surfaces using hammer and chisels. A compass and clinometer were used to record the geometry of the sampled surfaces and the surrounding topography, and locations and altitudes were determined with a hand-held GPS unit cross-referenced to a 1:25,000 topographic map (Table 1).

All samples were crushed and sieved to 250–500 μm and preparation for ^{10}Be analysis followed the procedures described in Wilson *et al.* (2008), as modified in Glasser *et al.* (2009). Measurement of ^{10}Be by accelerator mass spectrometry (AMS) is described in detail by Xu *et al.* (2010). The $^{10}\text{Be}/^9\text{Be}$ isotope ratios were normalized to the National Institute of Standards and Technology Standard Reference Material corresponding to a $^{10}\text{Be}/^9\text{Be}$ nominal value of 2.79×10^{-11} for a ^{10}Be half-life of 1.36 Ma (Nishiizumi *et al.*, 2007). This standard agrees with standards prepared by K. Nishiizumi, which were used as secondary standards. Procedural blank corrections were in the range 3–9.6%. The standard uncertainties of the cosmogenic nuclide concentrations include the AMS counting statistics and scatter uncertainties from sample and blank measurements, which includes the long-term AMS and chemical preparation uncertainties.

Exposure age calculation and results

The ^{10}Be surface exposure ages were calculated using the two methods adopted by the BRITICE-CHRONO project (D. Small *pers. comm.*, 2016). This enables the Nursery Knott ages to be related directly to results from other areas in northern England deriving from that project.

First, ages were determined using version 2.3 of the online calculators formerly known as CRONUS-Earth ^{10}Be – ^{26}Al exposure age calculators ([http://hess.ess.washington.edu/math/al be v23/al be multiple v23.php](http://hess.ess.washington.edu/math/al%20be%20multiple%20v23.php) Balco *et al.*, 2008) using the independently constrained Loch Lomond production rate (LLPR; 3.92 ± 0.18 atoms $\text{g}^{-1} \text{a}^{-1}$) (Fabel *et al.*, 2012). Production rates have not been determined for the Yorkshire Dales but are unlikely to differ significantly from the LLPR given the relatively short distance (~280km) between Loch Lomond (Scotland) and the Yorkshire Dales. In version 2.3 of the CRONUS-Earth age calculators a value of 4.0 ± 0.17 atoms $\text{g}^{-1} \text{a}^{-1}$ is used for the LLPR rather than 3.92 ± 0.18 atoms $\text{g}^{-1} \text{a}^{-1}$, but the resulting differences in age are not significant (D. Small *pers. comm.*, 2016). The LLPR was established from a geochronology provided by radiocarbon dating (MacLeod *et al.*, 2011). Exposure ages were based on the time-dependent Lm scaling (Lal, 1991; Stone, 2000) and assume 1mm ka^{-1} of post-depositional surface erosion. This erosion rate is in general use in TCN dating studies across a range of crystalline rock types (cf. André, 2002; Nicholson, 2009; Larsen *et al.*, 2012) and for this reason has been adopted here.

Second, exposure ages were calculated using the CRONUScalc program v2.0 (Marrero *et al.*, 2016) using the default global production rate of 3.92 atoms $\text{g}^{-1} \text{a}^{-1}$ for Sa scaling (Borchers *et al.*, 2016) and an erosion rate of 1mm ka^{-1} .

Sample code	AMS ID	^{10}Be (10^4 atoms g^{-1})	Blank correction (%)	Exposure Age ¹ (LLPR)	Exposure Age ² (CRONUScalc)
NK-01	SUERCb9963	10.602 ± 0.329	7.5	18.51 ± 1.01 (0.59)	18.4 ± 1.5 (0.6)
NK-02A	SUERCb9966	8.140 ± 0.275	9.6	14.84 ± 0.83 (0.51)	14.8 ± 1.2 (0.5)
NK-03	SUERCb9967	12.087 ± 0.391	6.7	21.31 ± 1.18 (0.71)	21.0 ± 1.7 (0.7)
NK-04	SUERCb9968	26.163 ± 0.574	3.0	46.75 ± 2.39 (1.08)	44.9 ± 3.6 (1.0)

¹ Exposure ages are based on the time-dependent Lm scaling (Lal, 1991; Stone, 2000) and assume 1mm ka^{-1} erosion.

² Exposure ages are based on SA scaling and assume 1mm ka^{-1} erosion. CRONUScalc reports ages to one decimal place.

Table 2: Cosmogenic (^{10}Be) data and surface exposure ages with total uncertainties at 1σ for the Nursery Knott samples. Analytical uncertainties (1σ) are in parentheses.




Figure 3A: The 2m-high boulder from which sample NK-01 was obtained at the arrowed location. Survey pole divisions are 20cm.
3B: The piece of rock taken for ^{10}Be surface exposure dating from the arrowed location in A. Scale bar is 30cm in length.
3C: The rear view of Figure 3A showing bedrock surface (plinth) from which the 2m-high boulder (left) was either pushed or dragged. Arrow indicates location of sample NK-02A.
3D: Position of sample NK-02A in the centre of the bedrock plinth.
3E: Boulder and position of sample NK-03 (arrowed).
3F: Boulder and position of sample NK-04 (arrowed).

Both production rates agree within 1σ uncertainties with the range of production rates determined for other high latitude sites in the northern hemisphere (Small and Fabel, 2016). The cosmogenic (^{10}Be) data and exposure ages with uncertainties for each method of calculation applied are given in Table 2. In the following section LLPR ages are reported first followed by CRONUScalc ages.

One boulder sample (NK-04) returned ages of 46.75 ± 2.39 ka and 44.9 ± 3.6 ka, pre-dating the LGM. It is inferred that insufficient erosion of the boulder occurred during glacier transport and that the sample is compromised by the effects of nuclide inheritance. The bedrock sample (NK-02A) returned ages of 14.84 ± 0.83 ka and 14.8 ± 1.2 ka, spanning the terminal stages of the Late Devensian Dimlington Stadial (~ 28 – 15 ka) and the Lateglacial Interstadial (14.7 – 12.9 ka) at 1σ uncertainty, and the Younger Dryas Stadial (12.9 – 11.7 ka) at 2σ uncertainty,

and is significantly younger than previously reported ages for deglaciation of the Yorkshire Dales. Therefore, neither of these samples provide close limiting constraints on the loss of glacial ice cover. Further consideration of these ages is provided below. The other two boulder samples yielded ages of 18.51 ± 1.01 ka and 18.4 ± 1.5 ka (NK-01), and 21.31 ± 1.18 ka and 21.0 ± 1.7 ka (NK-03), consistent within 2σ uncertainties with the range of ages previously obtained for deglaciation of the Yorkshire Dales and other parts of northern England (Wilson *et al.*, 2012, 2013). However, NK-01 and NK-03 ages determined by use of the same production rate do not overlap within their 2σ analytical uncertainties (used for comparing samples from the same geological context) and have reduced Chi-square (χ^2_R) values of ~ 8 – 9.5 , indicating that they do not constitute a single population, for which a $\chi^2_R < 1$ is required.

Discussion

The ^{10}Be ages of the three boulders (NK-01, -03 and -04) indicate that the Stump Cross area was glaciated during the LGM, even though NK-04 has yielded a substantially earlier age than NK-01 and NK-03 (Table 2). Nevertheless, the pre-LGM age of NK-04 is useful in that it indicates that boulder transport and deposition must post-date its exposure ages of ~47–45 ka, and the LGM is the only glacial event in which the boulder could have been moved to its present location. Similar reasoning has been applied to explain boulders with ‘old’ ages from contexts in which LGM ice was clearly present (e.g. Fabel *et al.*, 2012; Ballantyne and Stone, 2015). Therefore, these three ages provide unequivocal evidence for the presence of glacier ice on the Wharfe–Nidd interfluvium, rather than the area having been above or beyond the influence of the last ice sheet (cf. Raistrick, 1931; Tillotson, 1934; Smith, 1977; Catt, 1991; Baker *et al.*, 1996; O’Connor and Lord, 2013).

Sample NK-02A is problematical in that it returned an age that is younger than expected given its position on the bedrock plinth from which boulder NK-01 was pushed or dragged. We hypothesize that following removal of the boulder the plinth was shielded from cosmic rays, for an unknown period, by a cover of glacial sediment and/or peat that has since been eroded. Walker (1956) noted that the erratic boulders around Nursery Knott are embedded in ‘glacial clay’; therefore, a former sediment cover across the plinth is not an unreasonable proposition.

Ages from boulder samples NK-01 and NK-03 are the only two that provide close limiting age constraints for deglaciation of this part of the Wharfe–Nidd interfluvium. Depending on the production rate applied and the resulting ages this event is defined as having occurred between 18.51 ± 1.01 ka and 21.31 ± 1.18 ka (LLPR) or between 18.4 ± 1.5 ka and 21.0 ± 1.7 ka (CRONUScal). Because these sample ages cannot be considered as deriving from the same statistical population they cannot be averaged to produce a best estimate age for deglaciation, and given that the relative accuracy of the LLPR and CRONUScal methods is unknown it is necessary to consider the merits of the age determinations for both boulders.

Several reconstructions depicting the retreat pattern of the last British Ice Sheet, or parts thereof, based on mapping, geomorphology, sedimentology, geochronology and physical numerical evidence have been produced in recent years (Evans *et al.*, 2009; Hubbard *et al.*, 2009; Clark *et al.*, 2012; Livingstone *et al.*, 2012, 2015; Chiverrell *et al.*, 2013; Bateman *et al.*, 2015; Hughes *et al.*, 2016). Each indicates a complex pattern of ice-sheet behaviour but there is a broad consensus that the Stump Cross area of the Yorkshire Dales remained ice covered until ~18 ka, after which there was a rapid loss of ice with the area becoming ice-free by ~17 ka. The exposure ages from boulder sample NK-01 lie closest to this proposed interval of deglaciation and may therefore be preferred, especially as the mean age of three erratics at Norber, 33 km to the northwest, is 17.9 ± 1 ka (Wilson *et al.*, 2012). Within 2σ total uncertainties the CRONUScal age for NK-03 also overlaps with the 18–17 ka period. In contrast, the NK-03 LLPR-derived age does not overlap with this interval suggesting that the sample may be compromised by slight nuclide inheritance.

The TCN ages for the Nursery Knott erratics are evidence of their deposition by glacial ice at ~18–17 ka following termination of the LGM and provide the first dated evidence for glaciation on this part of the Wharfe–Nidd interfluvium. The results resolve previous uncertainties relating to the age of the tills in the locality which can now be assigned to the LGM. The transport of substantial erratic blocks during the final phase of the LGM implies the presence of a glacier ice cover with the potential to re-activate the hydrology of relict stream passages in Stump Cross Caverns. However, there is no evidence from this cave system of the laminated clay units such as those reported from Victoria Cave, 25 km to the northwest, where they are

interpreted as being formed during hydrological rejuvenation associated with warm-based ice during middle and late Pleistocene glaciations (Lundberg *et al.*, 2010). Further work is clearly needed to investigate the effects of warm-based ice cover on the cave systems of the Yorkshire Dales, and to explain the variability in the sediments.

Conclusions

Being close to Stump Cross Caverns, the TCN results for the Nursery Knott boulder samples have provided an opportunity to compare surface and sub-surface dating evidence for deglaciation; the results from two boulders in this project and the previously published U–Th dates on speleothems are consistent within their inherent uncertainties and indicate deglaciation by ~17 ka. In the future more precise analysis by, for example, thermal ionization mass spectrometer U–Th dating of speleothems, has the potential to reduce these uncertainties still further.

Acknowledgements

We thank: the BCRA for funding towards the TCN analyses; Dr David Small (University of Glasgow) for calculating the ^{10}Be exposure ages using the LLPR; and an anonymous referee for spotting an error with a shielding value that resulted in an initial underestimate of one sample age. The samples were prepared at the NERC Cosmogenic Isotope Analysis Facility and measured at the Scottish Universities Environmental Research Centre AMS Laboratory.

References

- André, M-F, 2002. Rates of postglacial rock weathering on glacially scoured outcrops (Abisko-Riksgränsen area, 68 °N). *Geografiska Annaler*, Vol.84A, 139–150.
- Atkinson, T C, Lawson, T J, Smart, P L, Harmon, R S and Hess, J W, 1986. New data on speleothem deposition and palaeoclimate in Britain over the last forty thousand years. *Journal of Quaternary Science*, Vol.1, 67–72.
- Baker, A, Smart, P L and Edwards, R L, 1996. Mass spectrometric dating of flowstones from Stump Cross Caverns and Lancaster Hole, Yorkshire: palaeoclimate implications. *Journal of Quaternary Science*, Vol.11, 107–114.
- Balco, G, Stone, J O, Lifton, N A and Dunai, T J, 2008. A complete and easily accessible means of calculating surface exposure ages or erosion rates from ^{10}Be and ^{26}Al measurements. *Quaternary Geochronology*, Vol.3, 174–195.
- Ballantyne, C K and Stone, J O, 2015. Trimlines, blockfields and the vertical extent of the last ice sheet in southern Ireland. *Boreas*, Vol.44, 277–287.
- Bateman, M D, Evans, D J A, Buckland, P C, Connell, E R, Friend, R J, Hartmann, D, Moxon, H, Fairburn, W A, Panagiotakopulu, E and Ashurst, R A, 2015. Last glacial dynamics of the Vale of York and North Sea lobes of the British and Irish Ice Sheet. *Proceedings of the Geologists’ Association*, Vol.126, 712–730.
- Borchers, B, Marrero, S, Balco, G, Caffee, M, Goehring, B, Lifton, N, Nishiizumi, K, Phillips, F, Schaefer, J and Stone, J, 2016. Geological calibration of spallation production rates in the CRONUS-Earth project. *Quaternary Geochronology*, Vol.31, 188–198.
- Bradwell, T, Stoker, M S, Golledge, N R, Wilson, C K, Merritt, J W, Long, D, Everest, J D, Hestvik, O B, Stevenson, A G, Hubbard, A L, Finlayson, A G and Mathers, H E, 2008. The northern sector of the last British Ice Sheet: maximum extent and demise. *Earth-Science Reviews*, Vol.88, 207–226.
- Caseldine, C J, McGarry, S F, Baker, A, Hawksworth, C and Smart, P L, 2008. Late Quaternary speleothem pollen in the British Isles. *Journal of Quaternary Science*, Vol.23, 193–200.
- Catt, J A, 1991. Late Devensian glacial deposits and glaciations in eastern England and the adjoining offshore region. 61–68 in Ehlers, J, Gibbard, P L and Rose, J (eds), *Glacial deposits in Great Britain and Ireland*. [Amsterdam: Balkema.]
- Chiverrell, R C, Burke, M J and Thomas, G S P, 2016. Morphological and sedimentary responses to ice mass interaction during the last deglaciation. *Journal of Quaternary Science*, Vol.31, 265–280.
- Chiverrell, R C, Thrasher, I M, Thomas, G S P, Lang, A, Scourse, J D, van Landeghem, K J J, McCarroll, D, Clark, C D, Ó Cofaigh, C, Evans, D J A and Ballantyne, C K, 2013. Bayesian modelling of the retreat of the Irish Sea Ice Stream. *Journal of Quaternary Science*, Vol.28, 200–209.

- Clark, C D, Evans, D J A, Khatwa, A, Bradwell, T, Jordan, C J, Marsh, S H, Mitchell, W A and Bateman, M D, 2004. Map and GISA database of glacial landforms and features related to the last British Ice Sheet *Boreas*, Vol.33, 359–375.
- Clark, C D, Hughes, A L C, Greenwood, S L, Jordan, C and Sejrup, H P, 2012. Pattern and timing of the retreat of the last British-Irish Ice Sheet. *Quaternary Science Reviews*, Vol.44, 112–146.
- Clark, P U, Dyke, A S, Shakun, J D, Carlson, A E, Clark, J, Wohlfarth, B, Mitrovica, JX, Hostetler, S W and McCabe, A M, 2009. The Last Glacial Maximum. *Science*, Vol.325, 710–714.
- Cockburn, H A P and Summerfield, M A, 2004. Geomorphological applications of cosmogenic isotope analysis. *Progress in Physical Geography*, Vol.28, 1–42.
- Dunham, K C and Stubblefield, C J, 1945. The stratigraphy, structure and mineralization of the Greenhow mining area, Yorkshire. *Quarterly Journal of the Geological Society of London*, Vol.100, 209–268.
- Evans, D J A, Livingstone, S J, Vieli, A and Ó Cofaigh, C, 2009. The palaeoglaciology of the central sector of the British and Irish Ice Sheet: reconciling glacial geomorphology and preliminary ice-sheet modelling. *Quaternary Science Reviews*, Vol.28, 739–757.
- Fabel, D, Ballantyne, C K and Xu, S, 2012. Trimlines, blockfields, mountain-top erratics and the vertical dimensions of the last British-Irish Ice Sheet in NW Scotland. *Quaternary Science Reviews*, Vol.55, 91–102.
- Gilmour, M, Currant, A P, Jacobi, R M and Stringer, C, 2007. Recent TIMS dating results from British Late Pleistocene vertebrate faunal localities: context and interpretation. *Journal of Quaternary Science*, Vol.22, 793–800.
- Glasser, N F, Clemmens, S, Schnabel, C, Fenton, C R and McHargue, L, 2009. Tropical glacier advances in the Cordillera Blanca, Peru between 15 and 9.5 ka, from cosmogenic ^{10}Be dating. *Quaternary Science Reviews*, Vol.28, 3448–3458.
- Gosse, J C and Phillips, F M, 2001. Terrestrial in situ cosmogenic nuclides: theory and application. *Quaternary Science Reviews*, Vol.20, 1475–1560.
- Hubbard, A, Bradwell, T, Gollidge, N, Hall, A, Patton, H, Sugden, D, Cooper, R and Stoker, M, 2009. Dynamic cycles, ice streams and their impact on the extent, chronology and deglaciation of the British-Irish Ice Sheet. *Quaternary Science Reviews*, Vol.28, 758–776.
- Hughes, A L C, Gyllencreutz, R, Lohne, Ø S, Mangerud, J and Svendsen, J I, 2016. The last Eurasian ice sheets – a chronological database and time-slice reconstruction, DATED-1. *Boreas*, Vol.45, 1–45.
- Lal, D, 1991. Cosmic ray labelling of erosion surfaces: in situ nuclide production rates and erosion models. *Earth and Planetary Science Letters*, Vol.104, 424–439.
- Larsen, N K, Linge, H, Håkansson, L and Fabel, D, 2012. Investigating the last deglaciation of the Scandinavian Ice Sheet in southwest Sweden with ^{10}Be exposure dating. *Journal of Quaternary Science*, Vol.27, 211–220.
- Livingstone, S J, Evans, D J A, Ó Cofaigh, C, Davies, B J, Merritt, J W, Huddart, D, Mitchell, W A, Roberts, D H and Yorke, L, 2012. Glaciodynamics of the central sector of the last British-Irish Ice Sheet in northern England. *Earth-Science Reviews*, Vol.111, 25–55.
- Livingstone, S J, Roberts, D H, Davies, B J, Evans, D J A, Ó Cofaigh, C and Gheorghiu, D M, 2015. Late Devensian deglaciation of the Tyne Gap Palaeo-Ice Stream, northern England. *Journal of Quaternary Science*, Vol.30, 790–804.
- Lundberg, J, Lord, T C and Murphy, P J, 2010. Thermal ionization mass spectrometer U–Th dates on Pleistocene speleothems from Victoria Cave, North Yorkshire, UK: Implications for palaeoenvironment and stratigraphy over multiple glacial cycles. *Geosphere*, Vol.6, 379–395.
- MacLeod, A, Palmer, A, Lowe, J, Rose, J, Bryant, C, Merritt, J, 2011. Timing of glacier response to Younger Dryas climatic cooling in Scotland. *Global and Planetary Change*, Vol.79, 264–274.
- Marrero, S M, Phillips, F M, Borchers, B, Lifton, N, Aumer, R and Balco, G, 2016. Cosmogenic nuclide systematics and the CRONUScal program. *Quaternary Geochronology*, Vol.31, 160–187.
- McCarroll, D, Stone, J O, Ballantyne, C K, Scourse, J D, Fifield, L K, Evans, D J A and Hiemstra, J F, 2010. Exposure-age constraints on the extent, timing and rate of retreat of the last Irish Sea ice stream. *Quaternary Science Reviews*, Vol.29, 1844–1852.
- Nicholson, D T, 2009. Holocene microweathering rates and processes on ice-eroded bedrock, Rødal area, Hardangervidda, southern Norway. 29–49 in Knight, J and Harrison, S (eds), *Periglacial and Paraglacial Processes and Environments*. Geological Society, London, Special Publications 320.
- Nishiizumi, K, Imamura, M, Caffee, M W, Southon, J R, Finkel, R C and McAninch, J, 2007. Absolute calibration of ^{10}Be AMS standards. *Nuclear Instruments and Methods in Physics Research B*, Vol.258, 403–413.
- O'Connor, T and Lord, T, 2013. Cave palaeontology. 225–238 in Waltham, T and Lowe, D (eds), *Caves and Karst of the Yorkshire Dales*. [Buxton: British Cave Research Association.]
- Ó Cofaigh, C, Dunlop, P and Benetti, S, 2012. Marine geophysical evidence for Late Pleistocene ice sheet extent and recession off northwest Ireland. *Quaternary Science Reviews*, Vol.44, 147–159.
- Phillips, W M, 2001. A review of cosmogenic nuclide surface exposure dating: new challenges for Scottish geomorphology. *Scottish Geographical Journal*, Vol.117, 1–15.
- Raistrick, A, 1931. The glaciation of Wharfedale, Yorkshire. *Proceedings of the Yorkshire Geological Society*, Vol.22, 9–30.
- Scourse, J D and Furze, M F A, 2001. A critical review of the glaciomarine model for Irish Sea deglaciation: evidence from southern Britain, the Celtic shelf and adjacent continental slope. *Journal of Quaternary Science*, Vol.16, 419–434.
- Sejrup, H P, Hjelstuen, B O, Torbjørn Dahlgren, K I, Haflidason, H, Kuijpers, A, Nygård, A, Praeg, D, Stoker, M S and Vorren, T O, 2005. Pleistocene glacial history of the NW European continental margin. *Marine and Petroleum Geology*, Vol.22, 1111–1129.
- Small, D and Fabel, D, 2016. Was Scotland deglaciated during the Younger Dryas? *Quaternary Science Reviews*, Vol.145, 259–263.
- Smith, R T, 1977. Stump Cross. 10 in Catt, J A (Ed.), *Guidebook for Excursion C7: Yorkshire and Lincolnshire*. X INQUA Congress Excursion Guides. [Norwich: Geo Abstracts.]
- Stone, J O, 2000. Air pressure and cosmogenic isotope production. *Journal of Geophysical Research: Solid Earth*, Vol.105, 23753–23759.
- Sutcliffe, A S, Lord, T C, Harmon, R S, Ivanovich, M, Rae, A and Hess, J W, 1985. Wolverine in northern England at about 83,000 yr B.P.: faunal evidence for climatic change during Isotope Stage 5. *Quaternary Research*, Vol.24, 73–86.
- Telfer, M W, Wilson, P, Lord, T C and Vincent, P J, 2009. New constraints on the age of the last ice sheet glaciation in NW England using optically stimulated luminescence dating. *Journal of Quaternary Science*, Vol.24, 906–915.
- Tillotson, E, 1934. The glaciation of Nidderdale. *Proceedings of the Yorkshire Geological Society*, Vol.22, 215–228.
- Vincent, P J, Wilson, P, Lord, T C, Schnabel, C and Wilcken, K M, 2010. Cosmogenic isotope (^{36}Cl) surface exposure dating of the Norber erratics, Yorkshire Dales: further constraints on the timing of the LGM deglaciation in Britain. *Proceedings of the Geologists' Association*, Vol.121, 24–31.
- Walker, D, 1956. A site at Stump Cross, near Grassington, Yorkshire, and the age of the Pennine microlithic industry. *Proceedings of the Prehistoric Society*, Vol.22, 23–28.
- Wilson, P, Bentley, M J, Schnabel, C, Clark, R and Xu, S, 2008. Stone run (block stream) formation in the Falkland Islands over several cold stages, deduced from cosmogenic isotope (^{10}Be and ^{26}Al) surface exposure dating. *Journal of Quaternary Science*, Vol.23, 461–473.
- Wilson, P, Barrows, T T, Lord, T C and Vincent, P J, 2012. Surface lowering of limestone pavement as determined by cosmogenic (^{36}Cl) analysis. *Earth Surface Processes and Landforms*, Vol.37, 1518–1526.
- Wilson, P, Lord, T C and Rodés, Á, 2013. Deglaciation of the eastern Cumbria glaciokarst, northwest England, as determined by cosmogenic nuclide (^{10}Be) surface exposure dating, and the pattern and significance of subsequent environmental changes. *Cave and Karst Science*, Vol.40, 22–27.
- Wilson, P and Lord, T C, 2014. Towards a robust deglacial chronology for the northwest England sector of last British-Irish Ice Sheet. *North West Geography*, Vol.14, 1–11.
- Xu, S, Dougans, A B, Freeman, S P H T, Schnabel, C and Wilcken, K M, 2010. Improved ^{10}Be and ^{26}Al AMS with a 5 MV spectrometer. *Nuclear Instruments and Methods B*, Vol.268, 736–738.